



**ASSOCIATION BETWEEN STAVUDINE ADMINISTRATION WITH  
LIPODYSTROPHY AND DYSLIPIDEMIA AMONG HIV-INFECTED  
PATIENTS IN DR. KARIADI HOSPITAL SEMARANG**

**JOURNAL OF MEDIA MEDIKA MUDA**

**Submitted to Fulfill the Requirements to Achieve Bachelor of Medicine  
Degree**

**JACOB BUNYAMIN  
22010110120041**

**UNDERGRADUATE PROGRAM OF MEDICINE  
FACULTY OF MEDICINE  
DIPONEGORO UNIVERSITY  
2014**

## DECLARATION OF APPROVAL

### DECLARATION OF APPROVAL

#### ASSOCIATION BETWEEN STAVUDINE ADMINISTRATION WITH LIPODYSTROPHY AND DYSLIPIDEMIA AMONG HIV-INFECTED PATIENTS IN DR. KARIADI HOSPITAL

Proposed by:

**JACOB BUNYAMIN**

**22010110120041**

Has been approved by:

Semarang, 14<sup>th</sup> July 2014

Supervisor,



**DR. Dr. Muchlis A.U. Sofro, Sp.PD-KPTI, FINASIM**  
196303191989031004

Chief Reviewer,



**Dr. Endang Sri Lestari, Ph.D**  
196610161997022001

Reviewer,



**DR. Dr. Shofa Chasani, Sp.PD-KGH,**  
**FINASIM**  
195102051979011001

# **HUBUNGAN ANTARA PEMBERIAN STAVUDINE DENGAN LIPODISTROFI DAN DISLIPIDEMIA PADA PENDERITA INFEKSI HIV DI RSUP DR. KARIADI SEMARANG**

## **ABSTRAK**

Jacob Bunyamin<sup>1</sup>, Muchlis A.U. Sofro<sup>2</sup>

**Latar belakang:** Pemberian stavudin (d4T) berkontribusi terhadap penurunan angka kematian pada penderita HIV. Di lain pihak, stavudin diketahui menimbulkan efek samping yang serius seperti lipodistrofi dan dislipidemia. Kedua efek samping tersebut dapat meningkatkan resiko terjadinya komplikasi pada jantung dan pembuluh darah.

**Tujuan:** Penelitian ini bertujuan untuk mengetahui hubungan antara pemberian stavudin dengan lipodistrofi dan dislipidemia pada penderita HIV di RSUP Dr. Kariadi dan juga mengetahui faktor-faktor yang berhubungan dengan lipodistrofi dan dislipidemia.

**Metode:** Penelitian ini adalah penelitian observasional analitik dengan metode belah lintang. Empat puluh sampel dibagi menjadi kelompok stavudin (23 sampel) dan kelompok kontrol (17). Seluruh responden diperiksa profil lipid dan lipodistrofi oleh dokter jaga di klinik VCT RSUP Dr. Kariadi. Data yang diperoleh dianalisis menggunakan uji Chi-Square dan uji Fisher.

**Hasil:** Prevalensi lipodistrofi sebesar 21,7% dan dislipidemia 82,6% pada kelompok stavudin. Pemberian stavudin berhubungan signifikan dengan dislipidemia ( $p=0.008$ ) terutama pada kenaikan trigliserida dan penurunan kolesterol HDL ( $p=0.048$  dan  $p=0.009$ ). Pemberian stavudin tidak berhubungan signifikan dengan lipodistrofi ( $p=0.051$ ) walaupun stavudin berhubungan dengan lipodistrofi pada pantat dan wajah ( $p=0.026$  dan  $p=0.013$ ). Jenis kelamin perempuan berhubungan dengan lipodistrofi pada kelompok stavudin ( $p=0.014$ ). Jenis kelamin, umur, hitung CD4 dan durasi terapi tidak berhubungan dengan dislipidemia pada kelompok stavudin.

**Kesimpulan:** Pemberian stavudin berhubungan dengan dislipidemia pada penderita HIV di RSUP Dr. Kariadi dan tidak berhubungan dengan lipodistrofi.

**Kata kunci:** stavudin, lipodistrofi, dislipidemia, HIV

<sup>1</sup> Mahasiswa Fakultas Kedokteran Universitas Diponegoro

<sup>2</sup> Staf Pengajar Departemen Penyakit Dalam Fakultas Kedokteran Universitas Diponegoro

**ASSOCIATION BETWEEN STAVUDINE ADMINISTRATION WITH  
LIPODYSTROPHY AND DYSLIPIDEMIA AMONG HIV-INFECTED  
PATIENTS IN DR. KARIADI HOSPITAL SEMARANG**

**ABSTRACT**

*Jacob Bunyamin<sup>1</sup>, Muchlis A.U. Sofro<sup>2</sup>*

**Background:** *Stavudine (d4T) administration has contributed in decreasing mortality rate of HIV-infected patients. In the other hand, stavudine is known to cause serious side effects such as lipodystrophy and dyslipidemia. Both side effects are known to increase the risk of developing cardiovascular complications.*

**Aims:** *This study aimed to determine the association between stavudine administration with lipodystrophy and dyslipidemia among HIV-infected patients in Dr. Kariadi Hospital and also determine the factors associated with lipodystrophy and dyslipidemia.*

**Methods:** *This study was an observational analytic study using cross-sectional method. Forty samples were divided into stavudine group (23 samples) and control group (17). All respondents had their lipid profile observed and lipodystrophy assessed by attending physicians in VCT clinic Dr. Kariadi Hospital. Obtained data were analyzed using Chi-Square Test and Fisher's Exact Test.*

**Results:** *The prevalence of lipodystrophy was 21.7% and dyslipidemia was 82.6% in stavudine group. Stavudine administration was significantly associated with dyslipidemia ( $p=0.008$ ) especially in triglycerides elevation and HDL-c depletion ( $p=0.048$  and  $p=0.009$ ). Stavudine administration was not significantly associated with lipodystrophy ( $p=0.051$ ) although it was associated with lipoatrophy of buttock and face ( $p=0.026$  and  $p=0.013$ ). Being female was associated with lipodystrophy incidence in stavudine group ( $p=0.014$ ). Sex, age, CD4 count, and duration of treatment were not associated with dyslipidemia in stavudine group.*

**Conclusion:** *Stavudine administration was associated with dyslipidemia among HIV-infected patients in Dr. Kariadi Hospital and was not associated with lipodystrophy.*

**Keywords:** *stavudine, lipodystrophy, dyslipidemia, HIV*

<sup>1</sup> *Student of Faculty of Medicine Diponegoro University*

<sup>2</sup> *Teaching Staff of Department of Internal Medicine Faculty of Medicine Diponegoro University*

## **Background**

The development of anti retroviral therapy (ART) has reduced mortality among HIV/AIDS patients. Coincident with these advances, was described a lipodystrophy syndrome occurring in patients taking ART. The syndrome is characterized by body fat redistribution and metabolic abnormalities. The metabolic disturbances reported are elevated triglyceride, cholesterol and fasting blood glucose level. A study taken in Uganda showed that prolonged exposure to ART in children contributes to metabolic changes thus increasing the cardiovascular risk of adulthood. It is proven that there is a direct connection between plasma cholesterol level and the arteriosclerotic process, especially the correlation between aortic fatty streaks in young persons with elevated serum cholesterol level.<sup>1-3</sup>

Stavudine, a Nucleoside Reverse Transcriptase Inhibitor (NRTI), has the highest propensity for causing mitochondrial dysfunction, yet it is relatively inexpensive and often has better availability in the developing world. According to 2013 WHO Antiretroviral Guidelines, stavudine administration should be discontinued due to its metabolic toxicities on mitochondria. A drug like stavudine has high capacity of inhibiting mtDNA  $\gamma$ -polymerase, which will cause mtDNA depletion and lipodystrophy as a further effect. A study in Western India showed that the prevalence of lipodystrophy was 46.1% and lipoatrophy was significantly associated with stavudine administration. A study in Tanzania resulted that 73% patients taking antiretroviral therapy suffered from dyslipidemia which was worse in patients who took stavudine.<sup>4-8</sup>

In Dr. Kariadi Hospital Semarang, there is little data known about the association between stavudine administration to lipodystrophy and dyslipidemia in Dr. Kariadi Hospital Semarang. The aim of this study is to determine the association between stavudine administration with lipodystrophy and dyslipidemia among HIV infected patients in Dr. Kariadi Hospital Semarang and also their associated routine demographic informations.

## **Methods**

This study was an observational analytic study using cross-sectional method. This study was conducted on March-May 2014. A preliminary survey was conducted on November 2013 – February 2014. The study took place in the Voluntary Counseling and Testing (VCT) Clinic of Dr. Kariadi Hospital Semarang. Samples were 40 HIV-infected patients administered with stavudine containing regimens in Dr. Kariadi Hospital for stavudine group (23 patients) and HIV-infected patients administered with zidovudine, lamivudine, and nevirapine for control group (17 patients).

The diagnosis of lipodystrophy was taken from Lipodystrophy Severity Grading Score (LSGS) sheet by attending physician in VCT clinic. Routine demographic informations such as age, sex, CD4 count, and duration of therapy were taken from case record forms. Patients were asked to fast overnight for 9 hours for lipid profile test. Lipid profiles checked were TG, TC, HDL-c, and

LDL-c blood level. A 5 ml blood sample was drawn from the patient's median cubital vein. The blood samples were analyzed using automated analyzer in Central Laboratory Dr. Kariadi Hospital Semarang.

Chi-square test was used to determine the association between age, sex, current CD4 count, and duration of therapy with the incidence of lipodystrophy and dyslipidemia. Ineligible data for chi-square test was analyzed using Fischer's Exact Test. All of data collections and research was under permission of the Commission of Health Research Bioethics Faculty of Medicine Diponegoro University/Dr. Kariadi Hospital Semarang Indonesia.

## Results and Discussions

**Table 1.** Samples Characteristics

Characteristics	Stavudine Group n (%)	Control Group n (%)	P Value Lipodystrophy	P Value Dyslipidemia
Sex			<b>0.014</b>	0.671
Male	12 (52.2)	9 (52.9)		
Female	11 (47.8)	8 (47.1)		
Age (years)			0.545	0.329
Mean	35.5	33.8		
Range	24 – 58	22 – 50		
Duration of treatment (months)			0.673	0.517
Mean	30.7	30.8		
Range	9 – 121	6 – 72		
CD4 Count (cell/mm <sup>3</sup> )			0.128	0.614
Mean	234.6	264.2		
Range	4 – 597	96 – 512		
ART regimens				
d4T+3TC+ NVP	6 (26.1)	-		
d4T+3TC+EVF	15 (65.2)	-		
d4T+3TC+TDF	2 (8.7)	-		
ZDV+3TC+NVP	-	17 (100)		

There were 21.7% patients with lipodystrophy in stavudine group. None of the patients in the control group acquired lipodystrophy. The prevalence is lower compared to a study in India, which was 46.1% for lipodystrophy. Another study in Australia reported that lipodystrophy in HIV-infected patients' prevalence reached 53%, in which 55% reported both peripheral lipoatrophy and central

lipohypertrophy, 31% with peripheral lipoatrophy only and 14% had central lipohypertrophy only. In Ethiopia, the prevalence was 68.3%.<sup>7,9-10</sup>

Dyslipidemia accounted in 82.6% of patients in stavudine group. This percentage is twice higher from the control group which was 41.2%. The percentage of normal lipid profile in stavudine group was 17.4% while the control group's was 58.8%. A study in Uganda showed that the prevalence of stavudine-associated dyslipidemia was 91%. A research in Kenya resulted that the prevalence of dyslipidemia was 63.1% in patients administered with stavudine.<sup>11-12</sup>

**Table 2.** Association between Stavudine and Lipodystrophy

Variables	Stavudine Group	Control Group	P Value
	n (%)	n (%)	
Lipodystrophy	5 (21.7)	0 (0.0)	0.051
Absent	18 (78.3)	17 (100.0)	
Total	23 (100.0)	17 (100.0)	

**Table 3.** Fat Distribution Changes Overview

Variables	Stavudine group		Control group		P value
	Fat distribution changes	Absent	Fat distribution changes	Absent	
	n (%)	n (%)	n (%)	n (%)	
Lipoatrophy					
Upper arm	4 (17.4)	19 (82.6)	0 (0.0)	17 (100)	0.097
Thigh	5 (21.7)	18 (78.3)	0 (0.0)	17 (100)	0.051
Buttock	6 (26.1)	17 (73.9)	0 (0.0)	17 (100)	<b>0.026</b>
Face	7 (30.4)	16 (69.6)	0 (0.0)	17 (100)	<b>0.013</b>
Lipohypertrophy					
Abdomen	8 (34.8)	15 (65.2)	6 (35.3)	11 (64.7)	0.616
Breasts	3 (13.0)	20 (87.0)	2 (11.8)	15 (88.2)	0.646
Nuchae	4 (17.4)	19 (82.6)	0 (0.0)	17 (100)	0.097
Chin	1 (4.3)	22 (95.7)	0 (0.0)	17 (100)	0.575

The statistical analysis showed that the administration of stavudine did not significantly correlate to lipodystrophy incidences in HIV-infected patients. A study in Rwanda found that stavudine administration had significant correlation with lipodystrophy (p=0.019). Van Griensven et al. showed that stavudine had significant association with facial lipodystrophy (p=0.011), thigh atrophy (p=0.006), and buttock atrophy (p=0.009) compared to zidovudine.<sup>1,13</sup>

These differences were likely caused by the lack of lipodystrophy definition in clinical settings. This study used Lipodystrophy Severity Grading Score (LSGS) measured by physical examination by attending physician while another study were conducted using data from patients' self reports or DEXA although Carter et al. stated that the results of both methods were balanced. A study concluded that the prevalence of lipodystrophy would be different because of different definitions of lipodystrophy applied which may result in different statistical significance. The sample size of this study was also considered less compared to various studies in other countries.<sup>14</sup>

Sex was the only factor that significantly correlated with lipodystrophy in stavudine group ( $p=0.014$ ). This finding was consistent with several studies conducted in Rwanda and South Africa which stated that being female was a significant risk factor in developing lipodystrophy. Females were also reported to have more central lipohypertrophy ( $p<0.0001$ ), a pattern of lipodystrophy found most frequently in this study. Females were known to have impaired lipid profile due to the lack of 17-beta estradiol after menopause which increased LDL-c and decreased HDL-c levels, although the median age of this study was 35.5 years old.<sup>10,13,15</sup>

Meanwhile, age, duration of treatment and CD4 count did not have significant correlation with lipodystrophy in HIV-infected patients in Dr. Kariadi Hospital ( $p=0.545$ ,  $p=0.128$  and  $p=0.673$ ). These findings were different with a study in Rwanda which stated that duration of treatment  $\geq 18$  months and CD4 count  $\geq 150$  cell/mm<sup>3</sup> had significant correlation with lipodystrophy ( $p=0.020$  and  $p=0.014$ ) while age was not significantly correlated ( $p=0.328$ ). These differences may be caused by different sample size and different methods of diagnosing lipodystrophy.<sup>13,16</sup>

Although lipodystrophy is seen as cosmetic problem, the presence of lipodystrophy may suggest metabolic disturbance such as dyslipidemia, impaired glucose level, or hyperlactatemia. Clinically diagnosed lipodystrophy may become useful hints in detecting metabolic abnormalities in a resource limited setting. The presence of lipodystrophy may also cause stigma in HIV-infected patients and discrimination. Hadigan et al. found that among HIV-infected patients with lipodystrophy, the LDL-c particle exhibited a proatherogenic pattern.<sup>17</sup>



**Table 4.** Association between Stavudine and Dyslipidemia

Variables	Stavudine Group	Control Group	<i>P</i> Value
	<b>n (%)</b>	<b>n (%)</b>	
Dyslipidemia	19 (82.6)	7 (41.2)	<b>0.008</b>
Normal	4 (17.4)	10 (58.8)	
Total	23 (100.0)	17 (100.0)	

**Table 5.** Lipid Profile Changes Overview

Variables	Stavudine group		Control group		<i>P</i> Value
	<b>Increased</b>	<b>Normal</b>	<b>Increased</b>	<b>Normal</b>	
	<b>n (%)</b>	<b>n (%)</b>	<b>n (%)</b>	<b>n (%)</b>	
TC (mg/dL)	10 (43.5)	13 (56.5)	4 (23.5)	13 (76.5)	0.166
TG (mg/dL)	11 (47.8)	12 (52.2)	3 (17.6)	14 (82.4)	<b>0.048</b>
HDL(mg/dL) <sup>1</sup>	12 (52.2)	11 (47.8)	2 (11.8)	15 (88.2)	<b>0.009</b>
LDL (mg/dL)	8 (34.8)	15 (65.2)	2 (11.8)	15 (88.2)	0.096

<sup>1</sup> : Decreased HDL

Stavudine administration was significantly associated with dyslipidemia among HIV-infected patients in Dr. Kariadi Hospital. This result was consistent with a study taken in Uganda which also compared stavudine with zidovudine ( $p=0.001$ ).<sup>11</sup>

Stavudine was significantly associated with elevated TG level and decreased HDL-c level. Administration of stavudine was not significantly correlated with both TC and LDL-c increases. This finding corresponded with the fact that dyslipidemia in HIV-infected patients is mostly characterized by hypertriglyceridemia and low HDL-c concentrations. Stavudine was also associated with higher TG elevation ( $p<0.05$ ).<sup>18-19</sup>

This study found that sex, age, CD4 count, and duration of therapy had no significant association with dyslipidemia in stavudine group. This finding was similar with a study in Uganda which stated that sex, age, CD4 count, and duration of treatment had no significant association with dyslipidemia.<sup>11</sup>

Although being male and age  $\geq 35$  were considered as traditional risk factors of dyslipidemia, this finding was not seen in HIV-infected patients treated with stavudine in Dr. Kariadi Hospital. Confounding factors such as unhealthy diet, tobacco smoking and sedentary lifestyle may have role in impaired lipid profiles since most patients reported to do sport less than three times in a week and several patients claimed to have smoking history.

Peak body weight > 65 kg, family history of diabetes mellitus, family history of hypertension, and lipodystrophy were determined as risk factors for dyslipidemia in HIV-infected patients in Uganda. This study excluded patients with history of metabolic syndrome due to already impaired lipid profiles prior to stavudine administration.<sup>11</sup>

Elevated TG level is known as a risk factor of atherosclerosis which leads to heart attack and stroke. This impaired lipid profile may also cause pancreatic inflammation. The level HDL-c is a more potent risk factor for coronary heart disease compared to LDL-c level in blood. A study found that the risk of developing coronary heart disease decreases 2-3% for every 1 mg/dL increase of HDL-c. Pharmacological intervention and lifestyle modification should be promoted to patients with impaired lipid profiles. A reduction in dietary carbohydrate intake may improve an atherogenic lipid profile.<sup>20</sup>

It is recommended to switch stavudine with safer NRTI such as zidovudine (AZT) in limited resources area. Abacavir (ABC) and tenofovir (TDF) may be used if the situation is possible. Studies showed that either abacavir or tenofovir administration may improve increased lipid profiles caused by stavudine although abacavir did not reverse lipodystrophy effectively.<sup>21-22</sup>

Several limitations of this study were the sample size was not large compared to other studies conducted in different countries. We also could not determine the association between Body Mass Index (BMI) with lipodystrophy and dyslipidemia because the lack of data needed to assess BMI of HIV-infected patients in Dr. Kariadi Hospital Semarang.

## **Conclusions**

Stavudine administration was significantly associated with dyslipidemia among HIV-infected patients in Dr. Kariadi Hospital and was not associated with lipodystrophy. Sex was significantly associated with lipodystrophy in stavudine group while age, CD4 count, and duration of treatment were not. None of the routine demographic informations were significantly associated with dyslipidemia in stavudine group.

## **Suggestions**

Stavudine should be phased out as first-line drugs in HIV treatment in Dr. Kariadi Hospital Semarang due to its metabolic side effects such as dyslipidemia and lipodystrophy. Zidovudine, abacavir or tenofovir should be used to replace stavudine due to its ability to improve disturbed lipid profiles in patients with stavudine associated dyslipidemia.

## Acknowledgements

The author would like to thank DR. Dr. Muchlis A.U. Sofro, Sp.PD-KPTI, FINASIM, Dr. Endang Sri Lestari, PhD and DR. Dr. Shofa Chasani, Sp.PD-KGH, FINASIM for all suggestions and advices during the research period. The author also would like to thank all respondents who had participated on this research.

## References

1. Joly V, Flandre P, Meiffredy V, Leturque N, Harel M, Aboulker P, et al. Increased risk of lipoatrophy under stavudine in HIV-1-infected patients: results of a substudy from a comparative trial. *AIDS*. 2002;16: 2447-54.
2. Piloya T, Bakeera-Kitaka S, Kekitiinwa A, Moses RK. Lipodystrophy among HIV-infected children and adolescents on highly antiretroviral therapy in Uganda: a cross sectional study. *Journal of International AIDS Society*. 2012;15:174-4.
3. Longo D, Fauci A, Kasper D, Hauser S, Jameson J, Loscalzo J. *Harrison's Principles of Internal Medicine*, 18<sup>th</sup> edition. Philadelphia: McGraw-Hill; 2011: 1137-64.
4. Troll, GJ. Approach to dyslipidemia, lipodystrophy, and cardiovascular risk in patients with HIV infection. *Curr Atherosclerotic Repository*. 2011;13: 51-56.
5. World Health Organization. Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection. Geneva: World Health Organization; 2013: 27-35.
6. Villaroya F, Domingo P, Giralt M. Lipodystrophy associated with highly active retroviral therapy in HIV infection: the adipocyte as a target of antiretroviral-induced mitochondrial toxicity. *Trends Pharmacology Sciences*. 2005;26(2):88-93.
7. Pujari SN, Dravid A, Naik E, Bhagat S, Tash K, Nadler JP, et al. Lipodystrophy and dyslipidemia among patients taking first-line, world health organization recommended highly active antiretroviral therapy regimens in western india. *Acquired Immune Deficiency Syndrome*. 2005;39:199-202.
8. Liu E, Armstrong C, Spiegelman D, Chalamilla G, Njelekela M, Hawkins C, et al. First-line antiretroviral therapy and changes in lipid levels over 3 years among HIV-infected adults in Tanzania. *CID*. 2013;56:1820-27.
9. Miller J, Carr A, Emery S et al. HIV lipodystrophy: prevalence, severity, and correlates of risk in Australia. *HIV Med*. 2003;4(3):293-301.
10. Feleke Y, Fekade D, Mezegebu Y. Prevalence of highly active antiretroviral therapy associated metabolic abnormalities and lipodystrophy in HIV-infected patients. *Ethiop Med Journal*. 2012;50(3):221-30.
11. Omech B, Sempa J, Castelnovo B, Opio K, Otim M, Mayanja-Kizza H, et al. Prevalence of HIV-associated metabolic abnormalities among patients taking first line antiretroviral therapy in Uganda. *IRSN AIDS*. 2012;18(5): 57-62.

12. Manuthu, EM. Prevalence of dyslipidemia and dysglycemia in HIV-infected patients at the Kenyatta National Hospital. *EAMJ*. 2008;85(1):10-7.
13. Van Griensven J, De Naeyer L, Mushi T et al. High prevalence of lipodystrophy among patients on stavudine-containing first-line antiretroviral therapy regimens in Rwanda. *Elsevier*. 2007;101;793-8.
14. Carter VM, Hoy JF, Bailey M, Colman PG, Nyulasi I, Mijch AM. The prevalence of lipodystrophy in an ambulant HIV-infected population: it all depends on the definition. *HIV Med*. 2001;2(3):174-180.
15. Szafran H, Smielak-Korombel W. The role of estrogens in hormonal regulation of lipid metabolism in women. *Przeql Lek*. 1998;55(5):266-70.
16. Andany N, Raboud JM, Walmsley S et al. Ethnicity and gender differences in lipodystrophy of HIV-positive individuals taking antiretroviral therapy in Ontario, Canada. *HIV Clin Trials*. 2011;12(2):89-103.
17. Hadigan C, Mazza S, Crum D, Grinspoon S. Rosiglitazone increases small dense low-density lipoprotein concentration and decreases high-density lipoprotein particle size in HIV-infected patients. *AIDS*. 2007;21:2543-46.
18. Aragonés G, Alonso-Villaverde C, Pardo-Reche P et al. Antiretroviral treatment-induced dyslipidemia in HIV-infected patients is influenced by the *APOC3*-related rs10892151 polymorphism. *BMC Medical Genetics*. 2011;12;120-6.
19. Anastos K, Lu D, Shi Q et al. Association of serum lipid levels with HIV serostatus, specific antiretroviral agents, and treatment regimens. *J Acquir Immune Defic Syndr*. 2007;45(1);34-42.
20. Gordon DJ, Probstfield JL, Garrison RJ, et al. High-density lipoprotein cholesterol and cardiovascular disease: four prospective american studies. *Circulation*. 1989;79:8-15.
21. Fisac C, Fumero E, Crespo M et al. Metabolic benefits 24 months after replacing a protease inhibitor with abacavir, efavirenz or nevirapine. *AIDS*. 2005;19(9):17-25.
22. Schewe CK, Maserati R, Wassmer G, Adam A, Weitner L. Improved lipid profiles and maintenance of virologic control in heavily pretreated HIV-infected patients who switched from stavudine to tenofovir treatment. *CID*. 2006;42(1):145-7.